Author Index

Abe, M., see Kanai, M. 567(1991)415
Altay, C., see Kutlar, F. 567(1991)359
Altieri, I., see Zuccaro, P. 567(1991)485
Arimoto, H., see Kamizono, A. 567(1991)113
Aronson, A. L., see Voyksner, R. D. 567(1991)389
Bakri, D., see Jouve, J. 567(1991)331
Bastian, T. L., see Herring, V. L. 567(1991)221
Becher, M., see Saeed, K. 567(1991)185
Beksaç, S., see Kutlar, F. 567(1991)359
Belpaire, F. M., see Vermeulen, A. M. 567(1991)472
Betto, P., see Zuccaro, P. 567(1991)485
Blaschke, G., see Radler, S. 567(1991)229
Bleyzac, N., see Boulieu, R. 567(1991)480
Boulieu, R., Bleyzac, N. and Ferry, S. High-performance liquid chromatographic determination of ganciclovir in plasma 567(1991)480
Brier, G., see Kovach, P. M. 567(1991)129
Brinkman, U. A. T., see Van der Horst, F. A. L. 567(1991)161
Burri, B. J., Kutnink, M. A. and Neidlinger, T. R. Assay of human transthyretin-bound holo-retinol-binding protein with reversed-phase high-performance liquid chromatography 567(1991)369
Carlqvist, J., see Lanbeck-Vallén, K. 567(1991)121
Chabraoui, L., Mathian, B., Patricot, M. C. and Revol, A. Specific assay for unconjugated dehydroepiandrosterone in human plasma by capillary gas chromatography with electron-capture detection 567(1991)299
Chakravarthy, B. K., see Mangalan, S. 567(1991)498
Chan, K., see Walubo, A. 567(1991)261
Chemke, J., see Zamir, I. 567(1991)319
Chen, C.-F., see Tsai, T.-H. 567(1991)405
Costi, P., see Zecca, L. 567(1991)425
Dandekar, K. A., see Janiszewski, J. S. 567(1991)274
Davidson, V. P. and Traiger, G. J. Determination of *p*-methylthiobenzamide and *p*-methylthiobenzamide-S-oxide from rat plasma using solid-phase extraction and high-performance liquid chromatography 567(1991)213
Dennhardt, R., see Zimmermann, J. 567(1991)240
Dingler, E., see Lohmann, A. 567(1991)506
Diquet, B., see Van Der Meersch-Mougeot, V. 567(1991)441
Dixit, V. and Dixit, V. M. Solid-phase extraction of 11-nor-Δ-9-tetrahydrocannabinol-9-carboxylic acid from human urine with gas chromatographic–mass spectrometric confirmation 567(1991)81
Dixit, V. M., see Dixit, V. 567(1991)81
Eriksson, B.-M. and Wikström, M. Determination of vanilmandelic acid in urine by coupled-column liquid chromatography combining affinity to boronate and separation by anion exchange 567(1991)1
Ferrario, P., see Zecca, L. 567(1991)425
Ferry, S., see Boulieu, R. 567(1991)480
Fukamachi, M., see Kawaguchi, S. 567(1991)11
Fukuda, H., Yamatodani, A., Imamura, I., Maeyama, K., Watanabe, T. and Wada, H. High-performance liquid chromatographic determination of histamine N-methyltransferase activity 567(1991)459
Fuller, R. W., see Murray, S. 567(1991)289
Gandhi, T. P., see Mangalan, S. 567(1991)498
Garg, V. and Jusko, W. J. Simultaneous analysis of prednisone, prednisolone and their major hydroxylated metabolites in urine by high-performance liquid chromatography 567(1991)39
Garver, F. A., see Kutlar, F. 567(1991)359
Goldberg, J. I., see Sloley, B. D. 567(1991)49
Goto, J., Saisho, Y. and Nambara, T. Studies on steroids. CCLII. Separation and characterization of 3-oxobile acids in serum by high-performance liquid chromatography with fluorescence detection 567(1991)343
Gramm, H.-J., see Zimmermann, J. 567(1991)240
Grushka, E., see Zamir, I. 567(1991)319
Guattari, B. High-performance liquid chromatographic determination, with ultraviolet detection, of S-adenosyl-L-methionine and S-adenosyl-L-homocysteine in rat tissues and simultaneously of normetanephrine and metanephrine for phenylethanolamine-N-methyltransferase or catechol-O-methyltransferase activities 567(1991)254
Gurgey, A., see Kutlar, F. 567(1991)359
Hachey, D. L., see Roy, D. 567(1991)309
Hammes, W. and Weyhenmeyer, R. New assay method for the determination of vinpocetine in human plasma by gas chromatography–mass spectrometry without iransesterification caused by solvents: a reply. 567(1991)504

Heebner, E. T., see Holland, M. L. 567(1991)433
Herault, J., see Jouve, J. 567(1991)331
Hermansson, J., see Wiese, B. 567(1991)175
Herring, V. L., Bastian, T. L. and Lalonde, R. L.
Solid-phase extraction and direct high-performance liquid chromatographic determination of metoprolol enantiomers in plasma 567(1991)221
Hirachi, N., see Kawaguchi, S. 567(1991)11
Hirai, Y., see Tomokuni, K. 567(1991)65
Hiramatsu, K., see Okuda, T. 567(1991)141
Holland, M. L. and Heebner, E. T.
High-performance liquid chromatographic assay with ultraviolet detection for the determination of etoperidone and two active metabolites, 5-(1-hydroxyethyl)etoperidone and 1-(3-chlorophenyl)piperazine, in plasma 567(1991)433
Holthuis, J. J. M., see Van der Horst, F. A. L. 567(1991)161
Homandberg, G. A., see Kang, K. 567(1991)57
Homma, M., see Kanai, M. 567(1991)415
Huang, B. L., see Kiang, C.-H. 567(1991)195
Huang, S.-K., see Xu, D.-K. 567(1991)451
Huang, Y.-S., see Redden, P. R. 567(1991)21
Huisman, T. H. J., see Kutlar, F. 567(1991)359
Ichiba, M., see Tomokuni, K. 567(1991)65
Iga, T., see Yamamura, Y. 567(1991)151
Imamura, I., see Fukuda, H. 567(1991)459
Inotsume, N., see Kamizono, A. 567(1991)113
Iwata, Y., see Kanai, M. 567(1991)415
Janiszewski, J. S., Shukla, U. A., Kaul, S., Dandekar, K. A. and Pittman, K. A.
High-performance liquid chromatographic assay of α-(4-fluorophenyl)-4-(5-fluoro-2-pyrimidinyl)-1-piperazine butanol (BMY 14802), a potential antipsychotic drug, in monkey and rat plasma 567(1991)274
Jouve, J., Bakri, D., Herault, J. and Muh, J. P.
Determination of urinary vanillactic acid and plasma dihydroxyphenylalanine as markers of non-secreting neuroblastoma by high-performance liquid chromatography with electrochemical detection 567(1991)331
Jusko, W. J., see Garg, V. 567(1991)39
Kamahori, M., Taki, M., Watanabe, Y. and Miura, J.
Analysis of plasma catecholamines by high-performance liquid chromatography with fluorescence detection: simple sample preparation for pre-column fluorescence derivatization 567(1991)351
Kamizono, A., Inotsume, N., Miyamoto, K., Ueda, K., Miyakawa, T., Arimoto, H. and Nakano, M.
Determination of sultopride and tiapride in serum by gas chromatography using a surface ionisation detector 567(1991)113
Kanai, M., Kouno, Y., Homma, M., Yamada, K., Oka, K., Noguchi, M., Abe, M., Sakakura, K. and Iwata, Y.
Simple and accurate determination of methylpyrazines in biofluids using high-performance liquid chromatography 567(1991)415
Kang, K., Schwarz, D., Sky-Peck, H., Kuettner, K. E. and Homandberg, G. A.
Novel method of purification of human leukocytic elastase using adsorption on a high-performance liquid chromatography gel permeation column 567(1991)57
Kaul, S., see Janiszewski, J. S. 567(1991)274
Kawaguchi, S., Hirachi, N. and Fukamachi, M.
New analyser for the determination of urinary vanillylmandelic acid, homovanillic acid and creatinine 567(1991)11
Kawakami, J., see Yamamura, Y. 567(1991)151
Kawanishi, E., see Makino, Y. 567(1991)248
Khetarpal, V. K., Storbeck, L. S., Wells, D. and Meacham, Jr., R. H.
Method for separation and determination of lactone and hydroxy acid forms of a new HMG CoA reductase inhibitor (RG 12561) in plasma 567(1991)491
Kiang, C.-H., Nolan, T., Huang, B. L. and Lee, C. P.-P.
Determination of femtomole/milliliter concentrations of enprostil acid in human plasma using high-performance liquid chromatography–laser-induced fluorescence detection 567(1991)195
Kiefer, C. R., see Kutlar, F. 567(1991)359
Kimura, S., see Kitagawa, T. 567(1991)267
Kitagawa, T., Tsuboi, I., Kimura, S. and Sasamoto, Y.
Rapid method for preparing a β-glucan-specific sensitive fraction from Limulus (*Tachypleus tridentatus*) amebocyte lysate 567(1991)267
Kohriyama, K., see Okuda, T. 567(1991)141
Kotaki, H., see Yamamura, Y. 567(1991)151
Kouno, Y., see Kanai, M. 567(1991)415
Kovach, P. M., Lantz, R. J. and Brier, G.
High-performance liquid chromatographic determination of loracarbef, a potential metabolite, cefaclor and cephalexin in human plasma, serum and urine 567(1991)129
Kuettner, K. E., see Kang, K. 567(1991)57

Kutlar, F., Moscoso, H., Kiefer, C. R., Garver, F. A., Beksaç, S., Önderoglu, L., Gurgey, A., Altay, Ç. and Huisman, T. H. J.
Quantities of adult, fetal and embryonic globin chains in the blood of eighteen- to twenty-week-old human fetuses 567(1991)359
Kutnink, M. A., see Burri, B. J. 567(1991)369
Kyoyama, K., see Okuda, T. 567(1991)141
Lalonde, R. L., see Herring, V. L. 567(1991)221
Lanbeck-Vallén, K., Carlqvist, J. and Nordgren, T.
Determination of ampicillin in biological fluids by coupled-column liquid chromatography and post-column derivatization 567(1991)121
Langvardt, P. W. and Nolan, R. J.
Determination of styrene-7,8-oxide in whole rat blood by gas chromatography–mass spectrometry 567(1991)93
Lantz, R. J., see Kovach, P. M. 567(1991)129
Lee, B. L., Ong, H. Y. and Ong, C. N.
Determination of thiamine and its phosphate esters by gradient-elution high-performance liquid chromatography 567(1991)71
Lee, C. P.-P., see Kiang, C.-H. 567(1991)195
Liehr, J. G., see Roy, D. 567(1991)309
Liu, X.-Q., see Xu, D.-K. 567(1991)451
Lohmann, A. and Dingler, E.
New assay method for the determination of vinpocetine in human plasma by gas chromatography–mass spectrometry without transesterification caused by solvents. Reply to Hammes and Weyhenmeyer 567(1991)506
Maeyama, K., see Fukuda, H. 567(1991)459
Makino, Y. and Kawanishi, E.
High-performance liquid chromatographic separation of human apotransferrin and monoferric and diferric transferrins 567(1991)248
Mangalan, S., Patel, R. B., Gandhi, T. P. and Chakravarthy, B. K.
Detection and determination of free and plasma protein-bound astemizole by thin-layer chromatography: a useful technique for bioavailability studies 567(1991)498
Mannucci, C., see Triolo, A. 567(1991)105
Mathian, B., see Chabraoui, L. 567(1991)299
McGee, J. P., Palin, K. J., Shaw, P. N. and Potter, C.
High-performance liquid chromatographic analysis of trilostane and ketotrilostane in rat plasma 567(1991)282
Meacham, Jr., R. H., see Khetarpal, V. K. 567(1991)491
Miura, J., see Kamahori, M. 567(1991)351
Miyakawa, T., see Kamizono, A. 567(1991)113
Miyamoto, K., see Kamizono, A. 567(1991)113
Moscoso, H., see Kutlar, F. 567(1991)359
Muh, J. P., see Jouve, J. 567(1991)331
Murray, S., Wellings, R., Taylor, I. K., Fuller, R. W. and Taylor, G. W.
Gas chromatographic–mass spectrometric assay to measure urinary N^{τ}-methylhistamine excretion in man 567(1991)289
Nakano, M., see Kamizono, A. 567(1991)113
Nambara, T., see Goto, J. 567(1991)343
Neidlinger, T. R., see Burri, B. J. 567(1991)369
Nishida, M., see Okuda, T. 567(1991)141
Noguchi, M., see Kanai, M. 567(1991)415
Nolan, R. J., see Langvardt, P. W. 567(1991)93
Nolan, T., see Kiang, C.-H. 567(1991)195
Nordgren, T., see Lanbeck-Vallén, K. 567(1991)121
Oka, K., see Kanai, M. 567(1991)415
Okuda, T., Nishida, M., Sameshima, I., Kyoyama, K., Hiramatsu, K., Takehara, Y. and Kohriyama, K.
Determination of atropine in biological specimens by high-performance liquid chromatography 567(1991)141
Önderoglu, L., see Kutlar, F. 567(1991)359
Ong, C. N., see Lee, B. L. 567(1991)71
Ong, H. Y., see Lee, B. L. 567(1991)71
Pacifici, R., see Zuccaro, P. 567(1991)485
Palin, K. J., see McGee, J. P. 567(1991)282
Patel, R. B., see Mangalan, S. 567(1991)498
Patricot, M. C., see Chabraoui, L. 567(1991)299
Perico, A., see Triolo, A. 567(1991)105
Pichini, S., see Zuccaro, P. 567(1991)485
Pini, L. A., see Zuccaro, P. 567(1991)485
Pittman, K. A., see Janiszewski, J. S. 567(1991)274
Post, M. H., see Van der Horst, F. A. L. 567(1991)161
Potter, C., see McGee, J. P. 567(1991)282
Radler, S. and Blaschke, G.
Enantioselective determination of dimethindene in urine after oral administration of racemic dimethindene 567(1991)229
Redden, P. R. and Huang, Y.-S.
Automated separation and quantitation of lipid fractions by high-performance liquid chromatography and mass detection 567(1991)21
Revol, A., see Chabraoui, L. 567(1991)299
Ricciarello, G., see Zuccaro, P. 567(1991)485
Rosseel, M. T., see Vermeulen, A. M. 567(1991)472

Roy, D., Hachey, D. L. and Liehr, J. G.
Determination of estradiol 2- and 4-hydroxylase activities by gas chromatography with electron-capture detection 567(1991)309
Saeed, K. and Becher, M.
On-line solid-phase extraction of piroxicam prior to its determination by high-performance liquid chromatography 567(1991)185
Saisho, Y., see Goto, J. 567(1991)343
Sakakura, K., see Kanai, M. 567(1991)415
Sameshima, I., see Okuda, T. 567(1991)141
Santa, T., see Yamamura, Y. 567(1991)151
Sasamoto, Y., see Kitagawa, T. 567(1991)267
Sawada, Y., see Yamamura, Y. 567(1991)151
Schwarz, D., see Kang, K. 567(1991)57
Shafiq-Ur-Rehman
Rapid isocratic method for the separation and quantification of major phospholipid classes by high-performance liquid chromatography 567(1991)29
Shaw, P. N., see McGee, J. P. 567(1991)282
Shukla, U. A., see Janiszewski, J. S. 567(1991)274
Sky-Peck, H., see Kang, K. 567(1991)57
Sloley, B. D. and Goldberg, J. I.
Determination of γ-glutamyl conjugates of monoamines by means of high-performance liquid chromatography with electrochemical detection and application to gastropod tissues 567(1991)49
Sternieri, E., see Zuccaro, P. 567(1991)485
Storbeck, L. S., see Khetarpal, V. K. 567(1991)491
Takehara, Y., see Okuda, T. 567(1991)141
Taki, M., see Kamahori, M. 567(1991)351
Taylor, G. W., see Murray, S. 567(1991)289
Taylor, I. K., see Murray, S. 567(1991)289
Teeuwsen, J., see Van der Horst, F. A. L. 567(1991)161
Tomokuni, K., Hirai, Y. and Ichiba, M.
Fluorimetric determination of hepatic δ-aminolevulinic acid synthase activity by high-performance liquid chromatography 567(1991)65
Traiger, G. J., see Davidson, V. P. 567(1991)213
Triolo, A., Mannucci, C. and Perico, A.
Gas chromatographic–mass spectrometric determination of two new antimycotic agents, 1-[(5-chloro-2-benzofuranyl)-(2-chlorophenyl)methyl]-1*H*-imidazole and 1-[(5-bromo-2-benzofuranyl)-phenylmethyl]-1*H*-imidazole, in rabbit plasma following topical administration: a preliminary comparison with bifonazole 567(1991)105
Tsai, T.-H. and Chen, C.-F.
High-performance liquid chromatographic determination of 18α-glycyrrhetinic acid and 18β-glycyrrhetinic acid in rat plasma: application to pharmacokinetic study 567(1991)405
Tsuboi, I., see Kitagawa, T. 567(1991)267
Tyczkowska, K. L., see Voyksner, R. D. 567(1991)389
Uchino, K., see Yamamura, Y. 567(1991)151
Uebis, V.
Determination of a new blood glucose-lowering agent in biological fluids by capillary gas chromatography using electron-capture detection 567(1991)465
Ueda, K., see Kamizono, A. 567(1991)113
Van der Horst, F. A. L., Van Opstal, M. A. J., Teeuwsen, J., Post, M. H., Holthuis, J. J. M. and Brinkman, U. A. T.
Comparative study on the determination of the anti-neoplastic drug teniposide in plasma using micellar liquid chromatography and surfactant-mediated plasma clean-up 567(1991)161
Van Der Meersch-Mougeot, V. and Diquet, B.
Sensitive one-step extraction procedure for column liquid chromatographic determination of fluvoxamine in human and rat plasma 567(1991)441
Van Opstal, M. A. J., see Van der Horst, F. A. L. 567(1991)161
Vermeulen, A. M., Rosseel, M. T. and Belpaire, F. M.
High-performance liquid chromatographic method for the simultaneous determination of *R*-(−)- and *S*-(+)-hexobarbital in rat plasma 567(1991)472
Voyksner, R. D., Tyczkowska, K. L. and Aronson, A. L.
Development of analytical methods for some penicillins in bovine milk by ion-paired chromatography and confirmation by thermospray mass spectrometry 567(1991)389
Wada, H., see Fukuda, H. 567(1991)459
Walubo, A., Chan, K. and Wong, C. L.
Simultaneous assay for isoniazid and hydrazine metabolite in plasma and cerebrospinal fluid in the rabbit 567(1991)261
Watanabe, T., see Fukuda, H. 567(1991)459
Watanabe, Y., see Kamahori, M. 567(1991)351
Wellings, R., see Murray, S. 567(1991)289
Wells, D., see Khetarpal, V. K. 567(1991)491
Weyhenmeyer, R., see Hammes, W. 567(1991)504

Wiese, B. and Hermansson, J.
Bioanalysis of diclofenac as its fluorescent carbazole acetic acid derivative by a post-column photoderivatization high-performance liquid chromatographic method 567(1991)175
Wikström, M., see Eriksson, B.-M. 567(1991)1
Wong, C. L., see Walubo, A. 567(1991)261
Xu, D.-K., Zhou, J.-H., Yuan, Y.-S., Liu, X.-Q. and Huang, S.-K.
High-performance liquid chromatographic assay for amiloride in plasma and urine 567(1991)451
Yamada, K., see Kanai, M. 567(1991)415
Yamamura, Y., Kawakami, J., Santa, T., Kotaki, H., Uchino, K., Sawada, Y. and Iga, T.
Selective high-performance liquid chromatographic method for the determination of glycyrrhizin and glycyrrhetic acid-3-O-glucuronide in biological fluids: application of ion-pair extraction and fluorescence labelling agent 567(1991)151
Yamatodani, A., see Fukuda, H. 567(1991)459
Yu, P. H., see Zuo, D. M. 567(1991)381
Yuan, Y.-S., see Xu, D.-K. 567(1991)451
Zamir, I., Grushka, E. and Chemke, J.
Separation and determination of saturated very-long-chain free fatty acids in plasma of patients with adrenoleukodystrophy using solid-phase extraction and high-performance liquid chromatographic analysis 567(1991)319
Zecca, L., Ferrario, P. and Costi, P.
Determination of diclofenac and its metabolites in plasma and cerebrospinal fluid by high-performance liquid chromatography with electrochemical detection 567(1991)425
Zhou, J.-H., see Xu, D.-K. 567(1991)451
Zimmermann, J., Dennhardt, R. and Granm, H.-J.
Measurement of plasma catecholamines by high-performance liquid chromatography with electrochemical detection in intensive care patients after dobutamine infusion 567(1991)240
Zuccaro, P., Altieri, I., Betto, P., Pacifici, R., Ricciarello, G., Pini, L. A., Sternieri, E. and Pichini, S.
Determination of naltrexone and 6β-naltrexol in plasma by high-performance liquid chromatography with coulometric detection 567(1991)485
Zuo, D. M. and Yu, P. H.
High-performance liquid chromatographic procedure for the simultaneous determination of aromatic L-amino acid decarboxylase activity towards 3,4-dihydroxyphenylalanine and 5-hydroxytryptophan 567(1991)381

Compound Index

S-Adenosyl-L-homocysteine
High-performance liquid chromatographic determination, with ultraviolet detection, of S-adenosyl-L-methionine and S-adenosyl-L-homocysteine in rat tissues and simultaneously of normetanephrine and metanephrine for phenylethanolamine-N-methyltransferase or catechol-O-methyltransferase activities 567(1991)254

S-Adenosyl-L-methionine
High-performance liquid chromatographic determination, with ultraviolet detection, of S-adenosyl-L-methionine and S-adenosyl-L-homocysteine in rat tissues and simultaneously of normetanephrine and metanephrine for phenylethanolamine-N-methyltransferase or catechol-O-methyltransferase activities 567(1991)254

Amiloride
High-performance liquid chromatographic assay for amiloride in plasma and urine 567(1991)451

L-Amino acid decarboxylase A
High-performance liquid chromatographic procedure for the simultaneous determination of aromatic L-amino acid decarboxylase activity towards 3,4-dihydroxyphenylalanine and 5-hydroxytryptophan 567(1991)381

δ-Aminolevulinic acid synthase
Fluorimetric determination of hepatic δ-aminolevulinic acid synthase activity by high-performance liquid chromatography 567(1991)65

Amoxicillin
Development of analytical methods for some penicillins in bovine milk by ion-paired chromatography and confirmation by thermospray mass spectrometry 567(1991)389

Ampicillin
Determination of ampicillin in biological fluids by coupled-column liquid chromatography and post-column derivatization 567(1991)121

Development of analytical methods for some penicillins in bovine milk by ion-paired chromatography and confirmation by thermospray mass spectrometry 567(1991)389

Apotransferrin
High-performance liquid chromatographic separation of human apotransferrin and monoferric and diferric transferrins 567(1991)248

Astemizole
Detection and determination of free and plasma protein-bound astemizole by thin-layer chromatography: a useful technique for bioavailability studies 567(1991)498

Atropine
Determination of atropine in biological specimens by high-performance liquid chromatography 567(1991)141

Biferric transferrin
High-performance liquid chromatographic separation of human apotransferrin and monoferric and diferric transferrins 567(1991)248

Bifonazole
Gas chromatographic–mass spectrometric determination of two new antimycotic agents, 1-[(5-chloro-2-benzofuranyl)(2-chlorophenyl)-methyl]-1*H*-imidazole and 1-[(5-bromo-2-benzofuranyl)phenylmethyl]-1*H*-imidazole, in rabbit plasma following topical administration: a preliminary comparison with bifonazole 567(1991)105

BMY-14802
High-performance liquid chromatographic assay of α-(4-fluorophenyl)-4-(5-fluoro-2-pyrimidinyl)-1-piperazine butanol (BMY 14802), a potential antipsychotic drug, in monkey and rat plasma 567(1991)274

1-[(5-Bromo-2-benzofuranyl)-phenylmethyl]-1*H*-imidazole
Gas chromatographic–mass spectrometric determination of two new antimycotic agents, 1-[(5-chloro-2-benzofuranyl)(2-chlorophenyl)-methyl]-1*H*-imidazole and 1-[(5-bromo-2-benzofuranyl)phenylmethyl]-1*H*-imidazole, in rabbit plasma following topical administration: a preliminary comparison with bifonazole 567(1991)105

Catecholamines
Measurement of plasma catecholamines by high-performance liquid chromatography with electrochemical detection in intensive care patients after dobutamine infusion 567(1991)240

High-performance liquid chromatographic determination, with ultraviolet detection, of S-adenosyl-L-methionine and S-adenosyl-L-homocysteine in rat tissues and simultaneously of normetanephrine and metanephrine for phenylethanolamine-N-methyltransferase or catechol-O-methyltransferase activities 567(1991)254

Analysis of plasma catecholamines by high-performance liquid chromatography with fluorescence detection: simple sample preparation for pre-column fluorescence derivatization 567(1991)351

Catechol-O-methyltransferase
High-performance liquid chromatographic determination, with ultraviolet detection, of S-adenosyl-L-methionine and S-adenosyl-L-homocysteine in rat tissues and simultaneously of normetanephrine and metanephrine for phenylethanolamine-N-methyltransferase or catechol-O-methyltransferase activities 567(1991)254

Cefaclor
High-performance liquid chromatographic determination of loracarbef, a potential metabolite, cefaclor and cephalexin in human plasma, serum and urine 567(1991)129

Cephalexin
High-performance liquid chromatographic determination of loracarbef, a potential metabolite, cefaclor and cephalexin in human plasma, serum and urine 567(1991)129

1-[(5-Chloro-2-benzofuranyl)(2-chlorophenyl)-methyl]-1*H*-imidazole
Gas chromatographic–mass spectrometric determination of two new antimycotic agents, 1-[(5-chloro-2-benzofuranyl)(2-chlorophenyl)-methyl]-1*H*-imidazole and 1-[(5-bromo-2-benzofuranyl)phenylmethyl]-1*H*-imidazole, in rabbit plasma following topical administration: a preliminary comparison with bifonazole 567(1991)105

5-(4-Chlorophenyl)-2-[(4-methylphenyl)-sulphonyl]-4-pentynoic acid
Determination of a new blood glucose-lowering agent in biological fluids by capillary gas chromatography using electron-capture detection 567(1991)465

1-(3-Chlorophenyl)piperazine
High-performance liquid chromatographic assay with ultraviolet detection for the determination of etoperidone and two active metabolites, 5-(1-hydroxyethyl)etoperidone and 1-(3-chlorophenyl)-piperazine, in plasma 567(1991)433

Cholesterol
Automated separation and quantitation of lipid fractions by high-performance liquid chromatography and mass detection 567(1991)21

Cholesteryl esters
Automated separation and quantitation of lipid fractions by high-performance liquid chromatography and mass detection 567(1991)21

Cloxacillin
Development of analytical methods for some penicillins in bovine milk by ion-paired chromatography and confirmation by thermospray mass spectrometry 567(1991)389

Creatinine
New analyser for the determination of urinary vanillylmandelic acid, homovanillic acid and creatinine 567(1991)11

Curdlan
Rapid method for preparing a β-glucan-specific sensitive fraction from Limulus (*Tachypleus tridentatus*) amebocyte lysate 567(1991)267

Dehydroepiandrosterone
Specific assay for unconjugated dehydroepiandrosterone in human plasma by capillary gas chromatography with electron-capture detection 567(1991)299

Diclofenac
Bioanalysis of diclofenac as its fluorescent carbazole acetic acid derivative by a post-column photoderivatization high-performance liquid chromatographic method 567(1991)175

Determination of diclofenac and its metabolites in plasma and cerebrospinal fluid by high-performance liquid chromatography with electrochemical detection 567(1991)425

Dihydroxyphenylalanine
Determination of urinary vanillactic acid and plasma dihydroxyphenylalanine as markers of non-secreting neuroblastoma by high-performance liquid chromatography with electrochemical detection 567(1991)331

3,4-Dihydroxyphenylalanine
High-performance liquid chromatographic procedure for the simultaneous determination of aromatic L-amino acid decarboxylase activity towards 3,4-dihydroxyphenylalanine and 5-hydroxytryptophan 567(1991)381

Dimethindene
Enantioselective determination of dimethindene in urine after oral administration of racemic dimethindene 567(1991)229

Dopamine
Determination of γ-glutamyl conjugates of monoamines by means of high-performance liquid chromatography with electrochemical detection and application to gastropod tissues 567(1991)49

Measurement of plasma catecholamines by high-performance liquid chromatography with electrochemical detection in intensive care patients after dobutamine infusion 567(1991)240

Analysis of plasma catecholamines by high-performance liquid chromatography with fluorescence detection: simple sample preparation for pre-column fluorescence derivatization 567(1991)351

Elastase
Novel method of purification of human leukocytic elastase using adsorption on a high-performance liquid chromatography gel permeation column 567(1991)57

Enprostil acid
Determination of femtomole/milliliter concentrations of enprostil acid in human plasma using high-performance liquid chromatography–laser-induced fluorescence detection 567(1991)195

Enzymes
Novel method of purification of human leukocytic elastase using adsorption on a high-performance liquid chromatography gel permeation column 567(1991)57

Fluorimetric determination of hepatic δ-aminolevulinic acid synthase activity by high-performance liquid chromatography 567(1991)65

High-performance liquid chromatographic determination, with ultraviolet detection, of S-adenosyl-L-methionine and S-adenosyl-L-homocysteine in rat tissues and simultaneously of normetanephrine and metanephrine for phenylethanolamine-N-methyltransferase or catechol-O-methyltransferase activities 567(1991)254

High-performance liquid chromatographic procedure for the simultaneous determination of aromatic L-amino acid decarboxylase activity towards 3,4-dihydroxyphenylalanine and 5-hydroxytryptophan 567(1991)381

High-performance liquid chromatographic determination of histamine N-methyltransferase activity 567(1991)459

Epinephrine
Measurement of plasma catecholamines by high-performance liquid chromatography with electrochemical detection in intensive care patients after dobutamine infusion 567(1991)240

Analysis of plasma catecholamines by high-performance liquid chromatography with fluorescence detection: simple sample preparation for pre-column fluorescence derivatization 567(1991)351

Estradiol
Determination of estradiol 2- and 4-hydroxylase activities by gas chromatography with electron-capture detection 567(1991)309

Etoperidone
High-performance liquid chromatographic assay with ultraviolet detection for the determination of etoperidone and two active metabolites, 5-(1-hydroxyethyl)etoperidone and 1-(3-chlorophenyl)-piperazine, in plasma 567(1991)433

Fatty acids
Separation and determination of saturated very-long-chain free fatty acids in plasma of patients with adrenoleukodystrophy using solid-phase extraction and high-performance liquid chromatographic analysis 567(1991)319

α-(4-Fluorophenyl)-4-(5-fluoro-2-pyrimidinyl)-1-piperazine butanol
High-performance liquid chromatographic assay of α-(4-fluorophenyl)-4-(5-fluoro-2-pyrimidinyl)-1-piperazine butanol (BMY 14802), a potential antipsychotic drug, in monkey and rat plasma 567(1991)274

Fluvoxamine
Sensitive one-step extraction procedure for column liquid chromatographic determination of fluvoxamine in human and rat plasma 567(1991)441

Free fatty acids
Separation and determination of saturated very-long-chain free fatty acids in plasma of patients with adrenoleukodystrophy using solid-phase extraction and high-performance liquid chromatographic analysis 567(1991)319

Ganciclovir
High-performance liquid chromatographic determination of ganciclovir in plasma 567(1991)480

Globin chains
Quantities of adult, fetal and embryonic globin chains in the blood of eighteen- to twenty-week-old human fetuses 567(1991)359

β-Glucans
Rapid method for preparing a β-glucan-specific sensitive fraction from Limulus (*Tachypleus tridentatus*) amebocyte lysate 567(1991)267

γ-Glutamyldopamine
Determination of γ-glutamyl conjugates of monoamines by means of high-performance liquid chromatography with electrochemical detection and application to gastropod tissues 567(1991)49

γ-Glutamyl-5-hydroxytryptamine
Determination of γ-glutamyl conjugates of monoamines by means of high-performance liquid chromatography with electrochemical detection and application to gastropod tissues 567(1991)49

γ-Glutamylmonoamines
Determination of γ-glutamyl conjugates of monoamines by means of high-performance liquid chromatography with electrochemical detection and application to gastropod tissues 567(1991)49

Glycyrrhetic acid-3-O-glucuronide
Selective high-performance liquid chromatographic method for the determination of glycyrrhizin and glycyrrhetic acid-3-O-glucuronide in biological fluids: application of ion-pair extraction and fluorescence labelling agent 567(1991)151

Glycyrrhetinic acid
High-performance liquid chromatographic determination of 18α-glycyrrhetinic acid and 18β-glycyrrhetinic acid in rat plasma: application to pharmacokinetic study 567(1991)405

Glycyrrhizin
Selective high-performance liquid chromatographic method for the determination of glycyrrhizin and glycyrrhetic acid-3-O-glucuronide in biological fluids: application of ion-pair extraction and fluorescence labelling agent 567(1991)151

Haemoglobin
Quantities of adult, fetal and embryonic globin chains in the blood of eighteen- to twenty-week-old human fetuses 567(1991)359

Hetacillin
Development of analytical methods for some penicillins in bovine milk by ion-paired chromatography and confirmation by thermospray mass spectrometry 567(1991)389

Hexobarbital enantiomers
High-performance liquid chromatographic method for the simultaneous determination of *R*-(−)- and *S*-(+)-hexobarbital in rat plasma 567(1991)472

Histamine N-methyltransferase
High-performance liquid chromatographic determination of histamine N-methyltransferase activity 567(1991)459

HMG CoA reductase inhibitor
Method for separation and determination of lactone and hydroxy acid forms of a new HMG CoA reductase inhibitor (RG 12561) in plasma 567(1991)491

Homovanillic acid
New analyser for the determination of urinary vanillylmandelic acid, homovanillic acid and creatinine 567(1991)11

Hydrazine
Simultaneous assay for isoniazid and hydrazine metabolite in plasma and cerebrospinal fluid in the rabbit 567(1991)261

Hydroxyestradiols
Determination of estradiol 2- and 4-hydroxylase activities by gas chromatography with electron-capture detection 567(1991)309

5-(1-Hydroxyethyl)etoperidone
High-performance liquid chromatographic assay with ultraviolet detection for the determination of etoperidone and two active metabolites, 5-(1-hydroxyethyl)etoperidone and 1-(3-chlorophenyl)-piperazine, in plasma 567(1991)433

Hydroxyprednisolones
Simultaneous analysis of prednisone, prednisolone and their major hydroxylated metabolites in urine by high-performance liquid chromatography 567(1991)39

20β-Hydroxyprednisone
Simultaneous analysis of prednisone, prednisolone and their major hydroxylated metabolites in urine by high-performance liquid chromatography 567(1991)39

5-Hydroxytryptamine
Determination of γ-glutamyl conjugates of monoamines by means of high-performance liquid chromatography with electrochemical detection and application to gastropod tissues 567(1991)49

5-Hydroxytryptophan
High-performance liquid chromatographic procedure for the simultaneous determination of aromatic L-amino acid decarboxylase activity towards 3,4-dihydroxyphenylalanine and 5-hydroxytryptophan 567(1991)381

Isoniazid
Simultaneous assay for isoniazid and hydrazine metabolite in plasma and cerebrospinal fluid in the rabbit 567(1991)261

Ketotrilostane
High-performance liquid chromatographic analysis of trilostane and ketotrilostane in rat plasma 567(1991)282

Laminaran
Rapid method for preparing a β-glucan-specific sensitive fraction from Limulus (*Tachypleus tridentatus*) amebocyte lysate 567(1991)267

Lichenan
Rapid method for preparing a β-glucan-specific sensitive fraction from Limulus (*Tachypleus tridentatus*) amebocyte lysate 567(1991)267

Lipids
Automated separation and quantitation of lipid fractions by high-performance liquid chromatography and mass detection 567(1991)21

Loracarbef
High-performance liquid chromatographic determination of loracarbef, a potential metabolite, cefaclor and cephalexin in human plasma, serum and urine 567(1991)129

Metanephrine
High-performance liquid chromatographic determination, with ultraviolet detection, of S-adenosyl-L-methionine and S-adenosyl-L-homocysteine in rat tissues and simultaneously of normetanephrine and metanephrine for phenylethanolamine-N-methyltransferase or catechol-O-methyltransferase activities 567(1991)254

N^1-Methylhistamine
Gas chromatographic–mass spectrometric assay to measure urinary N^1-methylhistamine excretion in man 567(1991)289

Methylpyrazines
Simple and accurate determination of methylpyrazines in biofluids using high-performance liquid chromatography 567(1991)415

***p*-Methylthiobenzamide**
Determination of *p*-methylthiobenzamide and *p*-methylthiobenzamide-S-oxide from rat plasma using solid-phase extraction and high-performance liquid chromatography 567(1991)213

***p*-Methylthiobenzamide-S-oxide**
Determination of *p*-methylthiobenzamide and *p*-methylthiobenzamide-S-oxide from rat plasma using solid-phase extraction and high-performance liquid chromatography 567(1991)213

Metoprolol
Solid-phase extraction and direct high-performance liquid chromatographic determination of metoprolol enantiomers in plasma 567(1991)221

Monoamines
Determination of γ-glutamyl conjugates of monoamines by means of high-performance liquid chromatography with electrochemical detection and application to gastropod tissues 567(1991)49

Monoferric transferrin
High-performance liquid chromatographic separation of human apotransferrin and monoferric and diferric transferrins 567(1991)248

6β-Naltrexol
Determination of naltrexone and 6β-naltrexol in plasma by high-performance liquid chromatography with coulometric detection 567(1991)485

Naltrexone
Determination of naltrexone and 6β-naltrexol in plasma by high-performance liquid chromatography with coulometric detection 567(1991)485

Neutrophil protease
Novel method of purification of human leukocytic elastase using adsorption on a high-performance liquid chromatography gel permeation column 567(1991)57

Norepinephrine
Measurement of plasma catecholamines by high-performance liquid chromatography with electrochemical detection in intensive care patients after dobutamine infusion 567(1991)240

Analysis of plasma catecholamines by high-performance liquid chromatography with fluorescence detection: simple sample preparation for pre-column fluorescence derivatization 567(1991)351

Normetanephrine
High-performance liquid chromatographic determination, with ultraviolet detection, of S-adenosyl-L-methionine and S-adenosyl-L-homocysteine in rat tissues and simultaneously of normetanephrine and metanephrine for phenylethanolamine-N-methyltransferase or catechol-O-methyltransferase activities 567(1991)254

11-Nor-Δ-9-tetrahydrocannabinol-9-carboxylic acid
Solid-phase extraction of 11-nor-Δ-9-tetrahydrocannabinol-9-carboxylic acid from human urine with gas chromatographic–mass spectrometric confirmation 567(1991)81

3-Oxobile acids
Studies on steroids. CCLII. Separation and characterization of 3-oxobile acids in serum by high-performance liquid chromatography with fluorescence detection 567(1991)343

Pachyman
Rapid method for preparing a β-glucan-specific sensitive fraction from Limulus (*Tachypleus tridentatus*) amebocyte lysate 567(1991)267

Penicillins
Development of analytical methods for some penicillins in bovine milk by ion-paired chromatography and confirmation by thermospray mass spectrometry 567(1991)389

Phenolethanolamine-N-methyltransferase
High-performance liquid chromatographic determination, with ultraviolet detection, of S-adenosyl-L-methionine and S-adenosyl-L-homocysteine in rat tissues and simultaneously of normetanephrine and metanephrine for phenylethanolamine-N-methyltransferase or catechol-O-methyltransferase activities 567(1991)254

Phosphatidylcholine
Rapid isocratic method for the separation and quantification of major phospholipid classes by high-performance liquid chromatography 567(1991)29

Phosphatidylethanolamine
Rapid isocratic method for the separation and quantification of major phospholipid classes by high-performance liquid chromatography 567(1991)29

Phosphatidylserine
Rapid isocratic method for the separation and quantification of major phospholipid classes by high-performance liquid chromatography 567(1991)29

Phospholipids
Automated separation and quantitation of lipid fractions by high-performance liquid chromatography and mass detection 567(1991)21

Rapid isocratic method for the separation and quantification of major phospholipid classes by high-performance liquid chromatography 567(1991)29

Piroxicam
On-line solid-phase extraction of piroxicam prior to its determination by high-performance liquid chromatography 567(1991)185

Prednisolone
Simultaneous analysis of prednisone, prednisolone and their major hydroxylated metabolites in urine by high-performance liquid chromatography 567(1991)39

Prednisone
Simultaneous analysis of prednisone, prednisolone and their major hydroxylated metabolites in urine by high-performance liquid chromatography 567(1991)39

Retinol-binding protein
Assay of human transthyretin-bound holo-retinol-binding protein with reversed-phase high-performance liquid chromatography 567(1991)369

RG 12561
Method for separation and determination of lactone and hydroxy acid forms of a new HMG CoA reductase inhibitor (RG 12561) in plasma 567(1991)491

Sphingomyelin
Rapid isocratic method for the separation and quantification of major phospholipid classes by high-performance liquid chromatography 567(1991)29

Steroids
Simultaneous analysis of prednisone, prednisolone and their major hydroxylated metabolites in urine by high-performance liquid chromatography 567(1991)39

Specific assay for unconjugated dehydroepiandrosterone in human plasma by capillary gas chromatography with electron-capture detection 567(1991)299

Studies on steroids. CCLII. Separation and characterization of 3-oxobile acids in serum by high-performance liquid chromatography with fluorescence detection 567(1991)343

Styrene-7,8-oxide
Determination of styrene-7,8-oxide in whole rat blood by gas chromatography–mass spectrometry 567(1991)93

Sultopride
Determination of sultopride and tiapride in serum by gas chromatography using a surface ionisation detector 567(1991)113

Teniposide
Comparative study on the determination of the anti-neoplastic drug teniposide in plasma using micellar liquid chromatography and surfactant-mediated plasma clean-up 567(1991)161

Thiamine
Determination of thiamine and its phosphate esters by gradient-elution high-performance liquid chromatography 567(1991)71

Thiamine phosphate esters
Determination of thiamine and its phosphate esters by gradient-elution high-performance liquid chromatography 567(1991)71

Tiapride
Determination of sultopride and tiapride in serum by gas chromatography using a surface ionisation detector 567(1991)113

Transferrins
High-performance liquid chromatographic separation of human apotransferrin and monoferric and diferric transferrins 567(1991)248

Transthyretin
Assay of human transthyretin-bound holo-retinol-binding protein with reversed-phase high-performance liquid chromatography 567(1991)369

Triacylglycerols
Automated separation and quantitation of lipid fractions by high-performance liquid chromatography and mass detection 567(1991)21

Trilostane
High-performance liquid chromatographic analysis of trilostane and ketotrilostane in rat plasma 567(1991)282

Vanillactic acid
Determination of urinary vanillactic acid and plasma dihydroxyphenylalanine as markers of non-secreting neuroblastoma by high-performance liquid chromatography with electrochemical detection 567(1991)331

Vanillylmandelic acid
New analyser for the determination of urinary vanillylmandelic acid, homovanillic acid and creatinine 567(1991)11

Vanilmandelic acid
Determination of vanilmandelic acid in urine by coupled-column liquid chromatography combining affinity to boronate and separation by anion exchange 567(1991)1

Very-long-chain free fatty acids
Separation and determination of saturated very-long-chain free fatty acids in plasma of patients with adrenoleukodystrophy using solid-phase extraction and high-performance liquid chromatographic analysis 567(1991)319

Vinpocetine
New assay method for the determination of vinpocetine in human plasma by gas chromatography–mass spectrometry without transesterification caused by solvents: a reply. 567(1991)504

New assay method for the determination of vinpocetine in human plasma by gas chromatography–mass spectrometry without transesterification caused by solvents. Reply to Hammes and Weyhenmeyer 567(1991)506

Vitamin A
Assay of human transthyretin-bound holo-retinol-binding protein with reversed-phase high-performance liquid chromatography 567(1991)369